

Gerald S. Bloomfield¹, **Maya G. Watanabe**², Sara M. McCallum³, Heather J. Ribaldo², Judith A. Aberg⁴, Aya Awwad³, Sarah M. Chu³, Judith S. Currier⁵, Carl J. Fichtenbaum⁶, Michael T. Lu³, Carlos D. Malvestutto⁷, Sophia Zhao³, Markella V. Zanni³, Steven Grinspoon³, Pamela S. Douglas¹ for REPRIEVE Investigators

¹Duke University School of Medicine, Durham, NC, USA, ²Harvard TH Chan School of Public Health, Boston, MA, USA, ³Massachusetts General Hospital, Boston, MA, USA, ⁴Icahn School of Medicine at Mt Sinai, New York, NY, USA, ⁵University of California Los Angeles, Los Angeles, CA, USA, ⁶University of Cincinnati, Cincinnati, OH, USA, ⁷The Ohio State University, Columbus, OH, USA

BACKGROUND

- People with HIV (PWH) have 2x higher risk of heart failure (HF).
- HF risk prediction is not well characterized in PWH due to lack of risk algorithm testing in this group.
- The AHA PREVENT HF 10-year risk score uses prior ASCVD risk equation components but performance has not been reported in PWH.
- REPRIEVE captured the prespecified and adjudicated outcome of HF in PWH allowing an opportunity to evaluate incidence of and factors associated with HF in a global cohort with low-moderate risk of cardiovascular disease.

METHODS

- Confirmed HF events included both hospitalized and non-hospitalized events:
 - Hospitalized events were independently adjudicated.
 - Non-hospitalized events were ascertained via the narrow standardized MedDRA query (SMQ) for heart failure.
- HF incidence rate (IR) (events/1000 person years) was described overall and by demographic, HF and HIV-specific factors.
- Average PREVENT HF 10-year risk scores were summarized overall and by risk factors.
 - Risk score components include sex, age, blood pressure, BMI, eGFR, diabetes, smoking, and anti-hypertensive medications.
- Expected number of HF events per participant per 1 year were calculated with 95% confidence intervals computed using 5,000 bootstrap samples.
- The PREVENT risk score was assessed using $\frac{\text{observed}}{\text{expected}}$ events ratio.

Among a global cohort of PWH with low-moderate ASCVD risk: age, sex, GBD region, race, obesity, and hypertension contributed to higher incidence of heart failure events

CONCLUSIONS

- In a global cohort of PWH with low-moderate ASCVD risk, the observed number of HF events with and without hospitalization was relatively low.
- Risk factors for HF events with and without hospitalization included older age, female sex, HIC, Black or African American race, high BMI, and hypertension (both controlled and uncontrolled).
- Expected events calculated using the AHA PREVENT HF risk score were similar to observed numbers of events overall and by participant characteristics.
- The PREVENT HF score provides a reasonable estimate of HF risk in PWH.

RESULTS

- Participant demographics are reported in **Table 1**.
- Overall IR of confirmed HF was 1.65/1000PY (median follow-up 5.6 years).
- Higher incidence of confirmed HF among: older age, female sex, high-income country (HIC) region, Black/African American race, current cigarette smoking, controlled/uncontrolled HTN (BP \leq / $>$ 140/90 mmHg), obesity (BMI \geq 30 kg/m²), and HIV-1 RNA \geq 400 copies/mL (**Figure 1A**).
- There was no apparent difference in HF IR by randomized treatment group.
- In HIC, the IR of confirmed HF was 2.32/1000PY compared to IR $<$ 0.8/1000PY in other GBD regions except Sub-Saharan Africa which was 1.67/1000PY.
- Among those in HIC, Black or African American participants had the highest IR of 3.68/1000PY compared to 1.46/1000PY for White participants.
- Among those with controlled hypertension, IR of HF was 3.31/1000PY compared to 1.97 and 1.13/1000PY for uncontrolled and no hypertension, respectively.
- There was higher incidence of HF in those with former or current smoking (2.09 and 2.35/1000PY) compared to never smoking.
- Most risk factors had observed/expected (O/E) events ratios close to or with confidence intervals spanning 1 with an overall O/E of 0.93 (95% CI: 0.71, 1.16) (**Figure 1B**).

Table 1. Participant Demographics

Demographics	N=7,769
Age, mean	50 (years)
Female Sex (%)	2419 (31%)
Non-White Race (%)	5065 (65%)
Current smoking (%)	1933 (25%)
Hypertension (HTN)* (%)	2780 (36%)
10-year ASCVD risk (%), median	4.5%
CD4, mean	659.8 cells/mm ³

* Includes HTN diagnosis or BP > 140/90 mmHg at entry




Figure 1. Incidence rates /1000PY (A) and ratio of observed/expected number of confirmed HF events (B). IRs were estimated using Poisson distribution. The IR x-axis is truncated at 5 with arrows indicating continuation of CI. Expected number of events are calculated per participant per 1 year with 95% CI computed using 5,000 bootstrap samples. †Due to low numbers of events, expected events were calculated by high- and non-high income GBD regions only.

Funding: REPRIEVE is supported through NIH grants U01HL123336, 1UG3HL164285, U01HL123339, and 1U24HL164284, the ACTG Network, Kowa Pharmaceuticals, Gilead Sciences, and ViiV Healthcare.

